



College of Veterinary Medicine

---

UNIVERSITY OF MINNESOTA

UNIVERSITY OF MINNESOTA RESEARCH PROJECT:

**TESTING THE ASSIST NPS PROGRAM**

**JUNE 18, 2014**

University of Minnesota | Department of Veterinary and Biomedical Sciences | College of Veterinary Medicine

---





**UNIVERSITY OF MINNESOTA RESEARCH PROJECT:**  
**TESTING THE ASSIST NPS PROGRAM**

A research study was conducted in the laboratory of Dr. Timothy Johnson at the University of Minnesota to test the impact of the ASSIST programmed approach on turkey performance and the turkey gastrointestinal microbiota.

---

**MATERIALS  
AND METHODS**

**EXPERIMENTAL DESIGN**

A total of 148 Hybrid Converter turkey toms were used in this study. Birds were acquired from a commercial source at day-of-hatch and randomly distributed into three dietary treatments with replicate pens (23 birds per pen, 6 pens total). The groups included: 1) poult receiving a standard corn-soy diet; 2) poult receiving a standard corn-soy diet supplemented with Quickstart plus mixed in feed (Assist Natural Products and Services, LLC, Lena, IL), Pro-Oxine supplemented continuously in drinking water (BioCide International, Norman, OK), and pre-treatment of litter shavings with Relentless Plus (Assist Natural Products and

Services, LLC, Lena, IL), all according to manufacturer's instructions; and 3) poult receiving a standard corn-soy diet, Quickstart plus supplemented continuously in drinking water (Assist Natural Products and Services, LLC, Lena, IL), Pro-Oxine supplemented continuously in drinking water (BioCide International, Norman, OK), and pre-treatment of litter shavings with Relentless Plus (Assist Natural Products and Services, LLC, Lena, IL), all according to manufacturer's instructions. Birds were followed from hatch through 42 days of age. Procedures regarding the use of the turkeys were approved by the Institutional Animal Care and Use Committee at the University of Minnesota, protocol 1309-30946A.

## SAMPLING

At days 3 and 7 of age, 8 birds per pen were randomly selected and euthanized. Each week following 7 days of age, 6 birds per pen were sampled. Staff were blinded to the experimental groups during bird postings. At euthanization, the following parameters were recorded: total bird weight, total intestinal weight, total intestinal length, and cecal score. Ileae from each bird were aseptically collected and processed for 16S rRNA microbiome profiling, as described below. From birds at 21 days of age, approximately 1 cm of the distal jejunum was collected from each bird and fixed in 10% formalin and embedded in paraffin. Villi heights and crypt depths were examined using formalin-fixed and paraffin-embedded samples under a light microscope at 100x, and expressed as an average of ten fields per sample.

---

## 16S rRNA BACTERIAL MICROBIOME PROFILING

DNA was extracted using a bead-beating procedure and the QIAmp<sup>®</sup> DNA Stool Kit (Qiagen, Valencia, CA) as previously described [1]. The V1-V3 hypervariable regions of the 16S rRNA gene were amplified in 25ul reactions containing 1X PCR buffer (containing 1.8 mM MgCl<sub>2</sub>), 0.2 mM each dNTP (Promega, Madison, WI), 0.4μM each primer (Integrated DNA Technologies, Coralville, IA), 1.25 U FastStart High Fidelity *Taq* polymerase (Roche,


Basel, Switzerland). Primers were designed for Illumina barcoding and sequencing as previously described [2]. Each forward and reverse primer contained a sample-specific sequence barcode. The PCR conditions used were an initial denaturation of 95°C for 2 min, followed by 25 cycles of 95°C for 30 sec, 60°C for 30 sec and 72°C for 30 sec; the amplification was completed with a final extension of 72°C for 7 min.

The PCR product was excised from a 1.5% gel and purified using the QIAquick Gel Extraction Kit following manufacturer's instructions (Qiagen). Sample DNA quality and quantity were assessed on a Bioanalyzer 2100 (Agilent, Palo Alto, CA) using a DNA-1000 lab chip. Sequencing was performed at the University of Minnesota using Illumina MiSeq paired-end 2X300 bp technology.

## DATA ANALYSIS

Following sequencing, sorting by barcode was performed to generate fastq files for each sample, which were quality-assessed and filtered prior to analysis. A de novo operational taxonomic unit (OTU) picking approach was used in QIIME [3] using uclust [4]. Potential chimeras were removed using ChimeraSlayer [4]. Approximately-maximum-likelihood phylogenetic trees were constructed using FastTree [5]. QIIME was also used for assessments of alpha diversity, beta diversity using Unifrac [6], and phylogenetic classifications using the RDP database [7,8]. Differential abundances of OTUs and other phylogenetic classifications were identified using METASTATS [9]. Construction of heatmaps was performed using the R statistical software [10].

### INTESTINAL LENGTH (cm)




## RESULTS

1

Sampling of intestinal length was performed by measuring length in cm from the proximal end of the small intestine through the ileocecal junction. No significant differences were observed in intestinal lengths at any of the timepoints between control and Assist-treated groups.


Control Dry Liquid

### INTESTINAL WEIGHT (g)



2

Similarly, intestinal weights were assessed at euthanization by weighing the same section of intestine used for intestinal lengths above. Again, no significant differences were observed in intestinal weights between control versus Assist-treated groups at any timepoint.




3

Total bird weights were also measured at euthanization in the control versus Assist-treated groups at each timepoint. At days 21 and 28 of age, both in-feed and in-water Assist treatment groups has significantly higher total body weights ( $P < 0.05$ ) than control groups. At day 35 of age, the in-feed treatment group weights were significantly higher ( $P < 0.05$ ) than the control group. At all timepoints after day 3, average weights were higher in Assist-treated groups than in the control groups.


4


Cumulative feed conversion rate (FCR) was assessed by calculating total feed consumed divided by total bird body weights for each experimental group. FCR values were 0.19-0.22 lower in Assist-treated groups compared to the control group.



5

Villi heights and crypt depths were also measured for individual birds from each experimental group at 21 days of age. For the in-feed treatment group, both villi heights and crypt depths were significantly larger ( $P < 0.05$ ) than for the control group; whereas, the crypt depths for the in-water treated group was significantly larger ( $P < 0.05$ ) than the control group at 21 days of age.





6

The total ileal bacteria for each individual bird in the experiment was assessed using 16S rRNA profiling. Based upon more than 1.5 million sequences generated (average >10,000 sequences per sample), a cladogram was generated to depict similarities between the ileal bacterial communities in each group / timepoint. From this analysis, ileal bacterial communities at days 21-42 of age were shifted in the Assist-treated groups compared to control groups of the same timepoints.

WEEK 1

Lactobacillus\_aviarius  
Lactobacillus\_salivarius  
Lactobacillus\_johnsonii

WEEK 2

Lactobacillus\_aviarius  
Clostridiales  
Lachnospiraceae  
Clostridium XIVa

WEEK 3

Lactobacillus\_crispatus  
Rothia

WEEK 4

Lactobacillus\_aviarius  
Lactobacillus\_johnsonii  
Erysipelotrichaceae  
Alistipes  
Faecalibacterium  
Lachnospiraceae  
Rothia  
Lactobacillus\_oris

WEEK 5

Lactobacillus  
Lactobacillus\_reuteri  
Lactobacillus\_reuteri  
Lactobacillus\_vaginalis  
Lactobacillus\_reuteri  
Lactobacillus\_reuteri  
Lactobacillus\_reuteri

WEEK 6

Lactobacillus\_crispatus  
Lactobacillus  
Clostridium\_XI

7

METASTATS was performed to determine members (bacterial species) of each community that were significantly enriched or depleted in the Assist-treated groups at each timepoint in the experiment. At weeks 4, 5, and 6, bacterial species that we have previously associated with enhanced performance in turkeys (*Lactobacillus aviarius*, *Lactobacillus johnsonii*, and *Clostridium group XI*) were enriched in the Assist-treated samples.

■ Enriched in the Assist-treated samples

## DISCUSSION

Our previous work has demonstrated that specific shifts in the turkey ileal microbiome correlates with enhanced performance in commercial turkeys [11]. Here, the use of Assist Natural Products programmatic approach was applied to commercial turkey toms over the course of 42 days to determine the impact of this approach on bird performance and the gut microbiota. This approach

included pre-treatment of litter with an amendment, continuous treatment of water with Pro-Oxine, and continuous administration of a prebiotic and probiotic mixture in feed or water. The results of this work indicate that this treatment significantly improves total bird weights, improves feed efficiency, and positively modulates the ileal bacterial microbiota. While it does not appear that significant

colonization of the bacterial species (*Bacillus* spp.) in the product are colonizing at high rates in the turkey gut, it is evident that this approach modulates the gut, including but gut development and microbiota maturation. It is likely that the effectiveness of these products are through indirect action on the environment and priming of the gut for colonization with beneficial commensal flora.

## REFERENCES CITED

1. Danzeisen JL, Kim HB, Isaacson RE, Tu ZJ, Johnson TJ (2011) Modulations of the chicken cecal microbiome and metagenome in response to anticoccidial and growth promoter treatment. *PLoS One* 6: e27949.
2. Bartram AK, Lynch MD, Stearns JC, Moreno-Hagelsieb G, Neufeld JD (2011) Generation of multimillion-sequence 16S rRNA gene libraries from complex microbial communities by assembling paired-end illumina reads. *Appl Environ Microbiol* 77: 3846-3852.
3. Caporaso JG, Kuczynski J, Stombaugh J, Bittinger K, Bushman FD, et al. (2010) QIIME allows analysis of high-throughput community sequencing data. *Nat Methods* 7: 335-336.
4. Edgar RC (2010) Search and clustering orders of magnitude faster than BLAST. *Bioinformatics* 26: 2460-2461.
5. Price MN, Dehal PS, Arkin AP (2010) FastTree 2--approximately maximum-likelihood trees for large alignments. *PLoS One* 5: e9490.
6. Lozupone C, Knight R (2005) UniFrac: a new phylogenetic method for comparing microbial communities. *Appl Environ Microbiol* 71: 8228-8235.
7. Wang Q, Garrity GM, Tiedje JM, Cole JR (2007) Naive Bayesian classifier for rapid assignment of rRNA sequences into the new bacterial taxonomy. *Appl Environ Microbiol* 73: 5261-5267.
8. Cole JR, Wang Q, Cardenas E, Fish J, Chai B, et al. (2009) The Ribosomal Database Project: improved alignments and new tools for rRNA analysis. *Nucleic Acids Res* 37: D141-145.
9. White JR, Nagarajan N, Pop M (2009) Statistical methods for detecting differentially abundant features in clinical metagenomic samples. *PLoS Comput Biol* 5: e1000352.
10. Dean CB, Nielsen JD (2007) Generalized linear mixed models: a review and some extensions. *Lifetime Data Anal* 13: 497-512.
11. Danzeisen JL, Calvert, A.J., Noll, S.L., McComb, B., Sherwood, J.S., Logue, C.M., Johnson, T.J. (2013) Succession of the turkey gastrointestinal bacterial microbiome related to weight gain. *PeerJ* 1: e237.

---

University of Minnesota | Department of Veterinary and Biomedical Sciences | College of Veterinary Medicine  
205 Veterinary Science | 1971 Commonwealth Ave. St. Paul, MN 55108 | Phone: 612.625.5255 Fax: 612.625.5203



Assist NPS | PO Box 784 | 218 Evergreen Dr. Lena, IL 61048 | Phone: 815.369.2195 Fax: 815.369.4001

---